

Article

## Effect of the Biofertilizer (*Bacillus Megaterium*) and the Addition of Yeast Spraying on the Vegetative Growths in Phosphorous Availability, Growth and Yield of Onions (*Allium Cepa* L.)

Hiba Mahdi Abdul-Hussein and Kareem U. Hassan

College of Agricultural Engineering, University of Baghdad, Iraq.

\*Correspondence: heba.mahdi1207a@coagri.uobaghdad.edu.iq

Available from: <http://dx.doi.org/10.21931/RB/CSS/2023.08.02.62>

**Abstract:** A field experiment was carried out in clay mixture soil according to the randomized complete block design with three replicates in one of the College of Agricultural Engineering Sciences / University of Baghdad fields. The study aims to identify the effect of the biofertilizer (*Bacillus megaterium*) and the addition of yeast on the availability of phosphorus, the yield, and the growth of onions after adding 50% of the fertilizer recommended for phosphorus. The study results showed the superiority of *Bacillus megaterium* treatment in the phosphorous availability, yield, vegetative growth and phosphatase enzyme activity (7.444 mg.kg<sup>-1</sup>, 0.540%, 0.298%, 35.521 µg), respectively. In comparison, the treatment of not adding *Bacillus megaterium* gave an average of (4.113 mg.kg<sup>-1</sup>, 0.363%, 0.198%, and 23.740 µg) respectively. Also, the results of the interaction between *Bacillus megaterium* and yeast showed the superiority of the treatment of adding yeast spraying to the vegetative part in the concentration of phosphorous in the soil. Besides, the yield, the vegetative part, and the activity of phosphatase enzyme (5.925 mg.kg<sup>-1</sup>, 0.480%, 0.292%, 32.483 µg), respectively, compared to the treatments of no yeast addition (5.630 mg.kg<sup>-1</sup>, 0.423%, 0.205%, and 26.778 µg).

**Keywords:** Biofertilizer, Yeast, Soil, Onions.

---

### Introduction

The element phosphorous is called (the key of life) because of its direct role in most of the vital processes inside the plant, as these processes cannot take place without it. It is likewise considered one of the essential nutrients for plant growth<sup>1</sup>. Phosphorous is included in the formation and division of living cells, the transfer of genetic traits, and the energy compound ATP and ADP, and it is considered one of the components of DNA and RNA; <sup>2</sup>. Furthermore, phosphorus participates in the decomposition of carbohydrates and substances that are the product of photosynthesis, and it also participates with proteins in the formation of cell membranes, such as the plasma membrane and the gap membrane. Plus, it has a role in increasing the yield by increasing the plant's resistance to some diseases and its role in the formation of seeds<sup>3</sup>. In the same role, it represents one of the essential nutrients for the growth of crops as it is a basis in vital physiological processes <sup>4</sup>. The world is moving towards clean agricultural technologies to reduce the pollution from chemical fertilizers as much as possible. Thus, the natural materials used, such as organic fertilizers and biofertilizers, are considered a suitable alternative to chemical fertilizers. <sup>5</sup>. *Bacillus* bacteria can dissolve phosphate compounds through their

ability to reduce the degree of soil reaction by producing organic acids such as citric and propionic acid. These acids can release phosphate through ion exchange with calcium, iron, or aluminum ions associated with phosphate<sup>6</sup>. Studies have indicated that yeast extract encourages cell division and thus leads to plant growth and an increase in leaf area, as well as the percentage of chlorophyll in the plant<sup>7</sup>. The addition of yeast extract to the vegetative growth affected the metabolic and vital activities of the plant. This extract had an encouraging effect that led to an increase in the photosynthetic pigments and enzymes responsible for this process that work to encourage growth in the plant.

Consequently, it increases the activity of the plant members, especially the leaves, because it produces photosynthesis products such as carotenoids and stores the products used later in forming the florets<sup>8</sup>. The use of yeast extract increases the characteristics of vegetative growth because it contains amino acids, nitrogen, and some mineral elements that stimulate growth and chlorophyll synthesis in the leaves. In addition, yeast is a natural source of cytokinins that have an essential role in cell division and increase their size<sup>9</sup>. Based on the foregoing, the study objective can be categorized as an evaluation of the role of *B. megaterium* inoculation in phosphorous availability in the plant rhizosphere by spraying yeast. Besides, assessment of spraying yeast to the vegetative growth in phosphorous availability, Studying and estimation of the activity of alkaline phosphatase enzyme in plant rhizosphere, growth, and yield of onions

## Materials and methods

### *Isolation, purification, and identification:*

10 soil samples were collected from the root zone (rhizosphere) of different plant hosts and geographical locations. Then, the samples were placed in sterile polyethylene bags and brought to the laboratory to conduct the process of *Bacillus megaterium* isolation. *Bacillus megaterium* was isolated in the microbiology laboratory in dishes containing a solid Pikovskaya culture medium with three replicates. The dishes were incubated upside down in the incubator at 30°C for (5) days. The growing colonies were examined to note that the colonies were surrounded by a transparent halo, which is evidence of the bacteria's ability to dissolve phosphate. The bacterial colonies were re-cultivated on the above solid culture media several times to ensure their ability to dissolve phosphate. Then, they were purified on the above culture media and then transferred to the slant medium for use in subsequent experiments. The following morphological, microscopic, and biochemical tests were carried out: colony color, colony shape, cell shape, cell arrangement, gram staining, catalase production, and oxidase production. Also, urease and starch decomposition, liquefaction of gelatin, motility, Voges-Proskauer, and the solubilization index were estimated.

### *Preparation of bacterial inoculum and activation of bacterial isolates*

100 ml of Nutrient broth was prepared and sterilized by autoclave at a temperature of 121°C and a pressure of 15 lb.in-2. After the medium was cooled, a portion of the isolated *Bacillus megaterium*, which had been previously isolated and identified, was taken by Loop. The liquid medium was inoculated with bacteria, and the inoculated medium was placed in the vibrating incubator for 48 hours. Then, the inoculum density was calculated, and a solution of gum Arabic was prepared at a concentration of 20% and sterilized with an autoclave at a temperature of 121 °C for a quarter of an hour before using it in the experiment.

### *Cultivation and inoculation*

The experiment was carried out at the Agricultural Research and Experiments Station of the College of Agricultural Engineering Sciences / University of Baghdad, where the field was divided into blocks according to the (RCBD) design using the drip irrigation system. The experiment included two factors; the first factor was the bio-fertilizer *Bacillus megaterium*, and the second factor was the yeast fungus, with three replicates. Onion sets (Texas White Crano) were planted on 7/9/2021 with 8 sets per furrow; the distance between one plant and another was 20 cm, and the number of plants reached 24 plants for each experimental unit. Before planting the onion sets, the biological inoculum of *Bacillus megaterium* was developed and prepared in the laboratory after washing the onion set with distilled water several times. Then, the bacterial inoculum was placed in a clean container. The Arabic gum was also placed at a rate of 20%, where the sets were placed in this gum, then the onion set was dipped in the biological inoculum several times for half an hour and placed on a clean cloth to dry it. After that, the set was planted in the field, where 5 g/l of yeast fungi were added in the form of spraying on the plant three sprays between each spray and another 15 days, 50% fertilizer recommendation was added to phosphorous.

Measured properties	Value	Unit	
Electrical conductivity EC 1:1	1.6	dS.m-1	
PH 1:1	7.28	---	
Available Nitrogen	20.5	Mg.kg-1 soil	
Available phosphorous	2.64		
Available potassium	132.41		
Soil separates	Sand	396	g.kg-1 soil
	Clay	344	
	Silt	260	
Texture	Clay loam		

**Table 1.** Chemical and physical soil analyses before planting.

### *Phosphatase activity test*

The activity of phosphatase enzyme in the soil before planting and after harvest was estimated, according to the (Tabatabai and Bremmer, 1965) method.

### **Results**

The results of biochemical tests indicated that they are starch-degrading bacteria, evidenced by the blue coloration of the medium, except for the colorless transparent halo around the bacterial growth area. Also, the fact that the medium of gelatin liquefaction remained in its liquid state when placed in the refrigerator indicated the ability of bacteria to liquefy gelatin and produce the enzyme gelatinase.

Isolate name and type. Test type	Eggplant B2	Alfalfa B5	Pepper B6
Colonial color	White	White	White
Colonial shape	Circular	Circular	Circular
Cell shape	Rod	Rod	Rod
Cell arrangement	Short strings	Short strings	Short strings
Gram stain	+	+	+
Catalyst production	+	+	+
Oxidase production	+	+	+
Decomposition of urease	-	-	-

<b>Decomposition of starch</b>	+	+	+
<b>Liquefying gelatin</b>	+	+	+
<b>Motility</b>	+	+	+
<b>Voges-Proskauer</b>	-	-	-

**Table 2. Results of diagnostic culture, morphological and biochemical tests for isolates of *B. megaterium*.**

*Results of determination of solubilization index (SI)*

The results showed the superiority of isolate (2), giving the highest solubilization index of 3.10 compared to the two isolates belonging to *B. megaterium*. In comparison, isolate (6) had the lowest solubilization index of 1.95.

Isolation number	Solubility	Plant host	SI
<b>B2</b>		Eggplant	3.10
<b>B5</b>		Alfalfa	2.89
<b>B6</b>		Pepper	1.95

**Table 3. Values of the solubilization index of *B. megaterium* isolates.**

*Effect of the biofertilizer (*Bacillus megaterium*) and the addition of yeast by spraying on the vegetative growth on the concentration of available phosphorus in the soil ( $\text{mg.kg}^{-1}$  soil)*

The results showed the superiority of the bio-fertilizer treatment in the phosphorous concentration in the soil with an average of ( $7.442$ )  $\text{mg.kg}^{-1}$  soil over the comparison treatment without *Bacillus megaterium* that was recorded ( $4.113$ )  $\text{mg.kg}^{-1}$  soil. The reason for this may be due to the ability of bacteria to dissolve insoluble phosphate compounds, as they have mechanisms in making phosphorous available for uptake by the plant to produce organic acids such as (succinic, lactic, and acetic acids) that increase the dissolution of insoluble phosphorous compounds. These results are consistent with <sup>14</sup> findings. However, the results of the interaction between biofertilizer and yeast showed that the treatment of adding yeast sprayed on the vegetative growth was superior to an average of ( $5.925$ )  $\text{mg.kg}^{-1}$  soil compared to the treatment of not adding yeast ( $5.630$ )  $\text{mg.kg}^{-1}$  soil. The reason for this is that yeast contains many substances that encourage better plant growth, which is reflected in an increase in the efficiency of phosphorous uptake <sup>15</sup>.

Biofertilizer	Yeast		Average
	Without addition	Addition by spraying	
<b>Without addition</b>	4.103	4.123	4.113
<b><i>Bacillus megaterium</i></b>	7.157	7.727	7.442
<b>Average</b>	5.630	5.925	
<b>LSD 0.05</b>	1.205	1.221	1.522

**Table 4. The effect of biofertilizer (*Bacillus megaterium*) and the addition of yeast by spraying on the vegetative growth on the concentration of available phosphorus in the soil ( $\text{mg.kg}^{-1}$  soil).**

*Effect of biofertilizer (*Bacillus megaterium*) and the addition of yeast by spraying to the vegetative growth on the concentration of phosphorous in the vegetative part %*

The results showed the superiority of the biofertilizer treatment in the phosphorous concentration of the vegetative part with an average of ( $0.298\%$ ) over the comparison treatment without *Bacillus megaterium*, which achieved ( $0.198\%$ ). This increase is attributed to the positive role of *B. megaterium*, which stimulates plant growth. Among them, it can increase the solubility of phosphate compounds in the soil by secreting organic acids such as oxalic, citric, gluconic, humic, acetic, and

formark, transforming phosphorous forms from unprepared forms to available plants. However, the interaction between the bio-fertilizer and yeast showed that the treatment of adding yeast sprayed on the vegetative part exceeded by an average of (0.292 %) compared to the treatment of not adding yeast achieved (0.205 %). The reason for this increase is attributed to the role of yeast in increasing the percentage of nitrogen and phosphorous in the vegetative growth <sup>17</sup>.

Biofertilizer	Yeast		Average
	Without addition	Addition by spraying	
Without addition	0.140	0.257	0.198
Bacillus megaterium	0.270	0.327	0.298
Average	0.205	0.292	
LSD 0.05	0.012	0.023	0.058

**Table 5. Effect of the biofertilizer (*Bacillus megaterium*) and the addition of yeast by spraying on the vegetative growth on the concentration of phosphorous in the vegetative part %.**

*Effect of the biofertilizer (*Bacillus megaterium*) and the addition of yeast by spraying on the vegetative growth on the concentration of phosphorous in yield %*

The results showed the superiority of the biofertilizer treatment in the phosphorous concentration in the yield with an average of (0.540 %) over the comparison treatment without *Bacillus megaterium* recorded (0.363 %). However, the interaction between the biofertilizer and yeast showed that adding yeast sprayed on the vegetative part exceeded by an average of (0.480 %) compared to the treatment of not adding yeast (0.423 %). This is due to the role of bread yeast in supplying the plant with essential mineral elements, some micro-elements, and vitamin B.

Biofertilizer	Yeast		Average
	Without addition	Addition by spraying	
Without addition	0.323	0.403	0.363
Bacillus megaterium	0.523	0.557	0.540
Average	0.423	0.480	
LSD 0.05	0.420	0.098	0.128

**Table 6. Effect of biofertilizer (*Bacillus megaterium*) and the addition of yeast by spraying on the vegetative growth on the concentration of phosphorous in yield %.**

*Effect of biofertilizer (*Bacillus megaterium*) and the addition of yeast by spraying on the vegetative growth in the phosphatase activity in the soil ( $\mu\text{g.g}^{-1} \text{ soil hour}^{-1}$ )*

The results showed the superiority of the biofertilizer treatment in the activity of the phosphatase enzyme, with an average of (35.521)  $\mu\text{g.g}^{-1} \text{ soil hour}^{-1}$ , over the comparison treatment without *Bacillus megaterium* (23.740)  $\mu\text{g.g}^{-1} \text{ soil hour}^{-1}$  even though the results of the interaction between biofertilizer and yeast showed the superiority of the yeast treatment spraying on the vegetative part with an average of (32.483)  $\mu\text{g.gm}^{-1} \text{ soil hour}^{-1}$  compared to the treatment without yeast addition (26,778)  $\mu\text{g.gm}^{-1} \text{ soil hour}^{-1}$ .

Biofertilizer	Yeast		Average
	Without addition	Addition by spraying	
Without addition	20.527	26.953	23.740
<i>Bacillus megaterium</i>	33.030	38.013	35.521
Average	26.778	32.483	
LSD 0.05	2.881	2.982	3.411

**Table 7.** The effect of the biofertilizer (*Bacillus megaterium*) and the addition of yeast by spraying on the vegetative growth on the phosphatase activity in soil  $\mu\text{g}\cdot\text{gm}^{-1}\text{ soil hour}^{-1}$ .

### Discussion

The superiority of the bacterial biofertilization treatment in the alkaline phosphatase activity is due to the ability of *B. megaterium* to secrete enzymes, especially the phosphatase enzyme, which increases the availability of phosphorous<sup>12</sup>. This discrepancy in the solubilization index may be due to its difference in genetic traits and secretion of enzymes, especially phosphatase enzymes and organic acids; this was mentioned<sup>12,13</sup>. The increase in phosphorus in the yield is attributed to the role of *B. megaterium* bacteria in the ability to dissolve phosphate compounds by producing organic acids and creating an appropriate PH reaction degree to increase phosphorus availability. Accordingly, the increase in its concentration in the soil and plants, reflected in the absorbed amount from the plant, led to an increase in plant growth<sup>18</sup>. Moreover, bacteria can move through the spread of growth outside the stabbing area, and the bacteria are oxidase-positive, evidenced by the change in color to dark violet and harmful for the Voges-Proskauer test for its inability to convert the color to red. These results are similar to what was found by several researchers in this domain<sup>10</sup>. However, these results present a good agreement with<sup>16</sup> findings. Finally, the role of yeast in increasing the percentage of nitrogen and phosphorous in vegetative growth is consistent with<sup>17</sup>.

### Conclusion

The microscopic culture tests showed that three phosphate-soluble isolates belonging to *B. megaterium* were obtained. The colonies were characterized by a circular shape surrounded by a round transparent halo with regular edges. The cell's shape was short rods, either singly or grouped, and they were gram-positive. Moreover, the examination showed a negative result of the urease test and the inability of the bacteria to change the medium color from yellow to pink, a catalase-positive bacteria that produces gas bubbles due to its ability to break hydrogen peroxide.

### References

1. Sharpley, A . Phosphorus availability. Handbook of Soil Science, 2000, D18-D30
2. Abu Dahi, Y. M.; Al-Younis, M. A. Plant Nutrition Index, Publishing House Directorate for Printing and Publishing, University of Baghdad, 1988.
3. Sanchez, C. A, Phosphorus. In-194 Barker, A. V. and Pilbeam D. J. (Eds) Handbook of plant nutrition. Taylor and Francis Crop LLC. Science. 2007 11(1):57-65.
4. Krishnaraj, P. U.; Dahale, S. Mineral phosphate solubilization: concepts and prospects in sustainable agriculture. In: *Proc Indian Natl Sci Acad.* 2014. p. 389-405.
5. Al-Wahaibi, M. b. H. Bacteria of the rhizosphere activating plant growth, *The Saudi Journal of Microbiological Sciences*, 2008. Volume (15), No. (3).
6. Ibraheem M W, Muhaimed A R, Mohammed Th. T. Leg cuts from Awaasi lambs fed a diet with varying levels of *Rhus coriaria* L., Physical dissection and chemical composition. *Revis Bionatura* 2022;7(4) 4. <http://dx.doi.org/10.21931/RB/2022.07.04.4>.

7. Devi, A. R.; Kotoky, R.; Pandey, P.; Sharma, G. D. Application of *Bacillus* spp. for sustainable cultivation of potato (*Solanum tuberosum* L.) and the benefits. In: *Bacilli and Agrobiotechnology*. Springer, Cham, 2016. p. 185-211.
8. El-Sherbeny, S. E.; Khalil, M.; Hessepn M. S. Growth and productivity of rue (*Ruta graveolens*) under different foliar fertilizers application. *Journal of Applied Sciences Research*, 2007, 3.5: 399-407.
9. El-Din, A. A. E., & Hendawy, S. F. Effect of dry yeast and compost tea on growth and oil content of *Borago officinalis* plant. *Res J Agric Biol Sci*, 2010, 6.4: 424-430.
10. Collee, J. G.; Miles, R. S.; Watt, B. Tests for The Identification of Bacteria, In *Practical Medical Microbiology* 14th ed., by J. Gerald Collee, Barrie, P., Marmion, Andrew, G., Fraser and Anthony Simmons, Churchill Livingstone, New York, 1996. p. 132-149.
11. Al-Hammadi, I. M. Identifying the effect of the growth of different plants on the microbial community in both the soil and the rhizosphere, University of Tripoli, Libya, *Aswat Journal of Agricultural Sciences*, 2014, 5: (45), 90-103.
12. Al-Obaidi, S. M. J. H. Effect of a combination of *Rhizobium phseoli*, *Bacillus megaterium* and *Bacillus mucilaginosus* with vermicompost on the NPK readiness in soil and in the growth and yield of beans (*Phaseolus vulgaris* L.), Master's thesis, College of Agricultural Engineering Sciences, University of Baghdad, Iraq. 2020.
13. Ahmed, F. A. Response of pot marigold (*Calendula officinalis* L.) plants to some rock phosphate sources and yeast. 5th Arab Hort. In: *Conf. Ismailia, Egypt*. 2001. p. 31-41.
14. Handhal, N. A.; Ahmaed, A. S. . A SURVEY STUDY TO ISOLATE SOME PATHOGENIC BACTERIA FOR COOKED RICE AT BAGHDAD CITY. *JLSAR* 2020, 1, 54-59.
15. Maysaloon W. Ibraheem, Abdulkhaliq A. Farhan, Sataa M. Salih, Th.T. Mohammed. Carcass characteristics of Awvasi lambs supplemented with Selenium and Vitamin D3. *Iranian Journal of Ichthyology*. 2022, Vol 9, No. 1, pp: 355-359..
16. Al-Dulaimi, A. F.; Farouk, F. J. Effect of spraying with yeast suspension, licorice extract and AMINO QVELANT-K on the chemical content of BLACK HAMBURG grape kernels, *Diyala Journal of Agricultural Sciences*, 2015, Volume (12), special issue of the fourth and first international scientific conference For Agricultural Research,. 15-16.
17. Abdel-Aziz, M. The Effect of Peanut Seed Size and Plant Spraying with *Saccharomyces Sereviciae* on Biological Yield and Some Indicators of Plant Productivity, *Syrian Journal of Agricultural Research*, 2018, 5(3): 125-134.
18. Rady, H. A. S.; Abdul-Ridha, H. A. Effect of locally produced biofertilizer from *Bacillus mucilaginosus* and *Bacillus subtilis* and imported biofertilizer on the growth and yield of *Solanum Tuberosum*. *Iraqi Journal of Desert Studies*, University of Anbar, Desert Study Center, 2018, 53- 65.

Received: May 15, 2023/ Accepted: June 10, 2023 / Published: June 15, 2023

Abdul-Hussein, H. M.; Hassan, K. U. Effect of the Biofertilizer (*Bacillus Megaterium*) and the Addition of Yeast Spraying on the Vegetative Growths in Phosphorous Availability, Growth and Yield of Onions (*Allium Cepa* L.). *Revis Bionatura* 2023;8 (2) 62. <http://dx.doi.org/10.21931/RB/CSS/2023.08.02.62>