

Microorganisms associated with bacterial wilt disease in *Dendrocalamus asper* (Giant Bamboo) from Ecuador

Jaime Morante-Carriel^{1,2*}, Marlon Vera-Carranza¹, Carlos Belezaca-Pinargote¹, Mercedes Carranza-Pa-tiño¹, Robinson J. Herrera-Feijoo^{1*}

¹Faculty of Agricultural and Forestry Sciences, Quevedo State Technical University, Av. Quito Km 1/2, vía a Santo Domingo de Los Tsáchilas. 120302, Quevedo, Ecuador

²Plant Proteomics and Functional Genomics Group, University of Alicante, C/ San Vicente del Raspeig, s/n. 03690, San Vicente del Raspeig, Alicante, Spain

* Correspondencia: rherreraf2@uteq.edu.ec; jmorante@uteq.edu.ec

Available from: <http://dx.doi.org/10.21931/RB/2024.09.01.32>

ABSTRACT

Dendrocalamus asper (Bamboo) is an essential component of forest ecosystems, whose health and survival are intrinsically linked to complex interactions with its microbiome. This study focuses on the pathogenic dynamics between the fungi *Arthrinium spp.* and *Erwinia spp.* Bacteria and their impact in the shoots wilting and culm rot in bamboo. Through humid chamber induction methods and selective cultivation techniques, pathogenic strains that significantly affect the regeneration and propagation of bamboo were identified and isolated. The investigation revealed a pathogenic synergy resulting in a high prevalence of diseases, 97% of the shoots evaluated showed signs of deterioration. Morphological identification challenged conventional techniques due to the variability of *Arthrinium spp.*—conidia, suggesting the need for more specific identification methods. The findings suggest that regulating soil pH and using beneficial biofilms could be promising strategies to mitigate bacterial infection. This study highlights the importance of sustainable management and in-depth knowledge of microbial ecology for integrated disease management in bamboo ecosystems. Understanding these microbial interactions is crucial for developing effective control strategies and long-term conservation of these vital ecosystems.

Keywords: *Dendrocalamus asper*, *Arthrinium spp.*, *Erwinia spp.*, bacterial wilt, disease management.

INTRODUCTION

Bamboo is vital in forest ecosystems, providing critical structure and support for microorganisms. Microorganisms, in particular, significantly influence the nutrition and health of bamboo. Studies have shown that bacteria such as *Flavobacterium*, *Bacillus*, and *Stenotrophomonas* are essential in absorbing nitrogen elements in moso bamboo, which could influence its flowering time¹. Bamboo planting can enrich soil fertility and increase tree species diversity in coastal ecosystems. In the bamboo forests of southeastern China, it has been observed that soil enzymatic and microbial activity is significantly altered². Microorganisms not only improve nutrient availability³. Microbial communities also protect against pathogens⁴, underscoring the importance of understanding microbial dynamics to sustain bamboo ecosystems.

Bamboo faces various diseases that can vary in prevalence and distribution worldwide, and microorganisms play a central role in these pathologies. Research has revealed that bamboo is an extensive reservoir of microorganisms, including fungi and bacteria, that should be investigated extensively, given its potential role in

plant growth and ecosystem functioning⁵. The invasion of native bamboo in subtropical forests can alter soil microbial communities. Disturbance affects litter decomposition and nutrient cycling. The threat of local forest ecosystems is affected due to microbial disturbance⁶. The findings highlight the need to carefully monitor bamboo planting and expansion to prevent adverse impacts on biodiversity and ecosystem health.

Adverse effects negatively correlate with soil pH and the abundance of stress, virulence, and sulfur-cycling genes⁷. Furthermore, specific genera such as *Arthrobacter*, *Pseudomonas*, *Acidobacteria* GP6, and *Pasteuria* may be potential biological control agents for bacterial wilt⁸. The findings underline the importance of understanding soil bacterial community structure for effectively managing bacterial wilt in bamboo.

The interaction between microorganisms and the plant response allows activating the defense mechanisms of *the Dendrocalamus asper* against microbial attacks crucial for survival. Research has shown that biofilm formation in the rhizosphere, observed in organic hydroponic systems, may be responsible for suppressing bacterial wilt⁹. Soil acidification aggravates the occurrence of bacterial wilt, suggesting that the regulation of soil acidification is a prerequisite and fundamental condition for disease control¹⁰. Previous findings are critical to developing management strategies that improve the physiological and biochemical responses of bamboo to bacterial infection, which could include modulation of soil pH and utilization of beneficial biofilms as part of an integrated approach to control diseases.

Microorganisms, including specific bacteria, play a crucial role in bacterial wilt that affects various plants, including *D. asper*. Studies have identified that strains of bacteria such as *Ralstonia solanacearum* vary considerably in Brazil, suggesting that host resistance to control bacterial wilt should be targeted primarily by region¹¹. Researchers have reported relevant findings on endophytic bacteria. Bacteria isolated from live oak trees in Texas showed the ability to inhibit the pathogen that causes oak wilt in vitro. Preinoculation with the bacteria *Pseudomonas denitrificans* 1-15 resulted in a 50% reduction in the incidence of diseased trees. Preinoculation decreased crown loss in affected trees by 17%¹². The findings highlight the importance of accurately identifying and classifying wilt-associated bacteria to develop effective control strategies.

D. asper possesses intrinsic defense mechanisms against microbial attacks, which include complex physiological and biochemical responses. Induction of resistance against diseases such as wilt caused by *Bursaphelenchus xylophilus* using selected pine endophytic bacteria is an example of how microorganisms can manage diseases by inducing systemic resistance in plants¹³. It has been observed that soil acidification can lead to outbreaks of bacterial wilt, underscoring the dynamic interaction between soil conditions and plant susceptibility to diseases¹⁴. The studies underscore the need to understand better plant responses to bacterial infection to improve disease management strategies.

The Bamboo *Dendrocalamus giganteus* is found in an environment where synergy between microorganisms can play a crucial role in its health and survival. Studies have shown that different species of tree-dwelling insects carry a variety of microorganisms, including pathogens, indicating their potential as vectors of plant diseases¹⁵. It has been observed that interactions between plant pathogens can lead to disease complexes that are more common than expected. Understanding these mechanisms can have essential implications in the epidemiology and management of plant diseases¹⁶. Therefore, it is plausible that a synergistic action between microorganisms could be present in pathogenic attacks on *D. giganteus* Munro. However, direct evidence specific to this bamboo species has yet to be established in the scientific literature.

The hypothesis that there is a synergy between different pathogenic microorganisms that contribute to bacterial wilt in *D. asper* in Ecuador is supported by the observation that interactions between plant attackers can be asymmetric, time-dependent and species-specific¹⁷. These complex interactions can influence the susceptibility of plants to diseases and their ability to resist or succumb to pathogen attacks. It has been reported that

plant pathogenic microorganisms can colonize insect tissues, revealing a network of interactions that affects substantial invaders¹⁸. The findings highlight the need to investigate further the possibility of microbial synergies in bacterial wilt of bamboo, particularly in regions such as Ecuador, where this disease can negatively impact bamboo production. This research aimed to identify and characterize the microorganisms associated with bacterial wilt diseases in *D. asper* (giant bamboo) from Ecuador.

MATERIALS AND METHODS

Plant material

The plant material was collected on the premises of the Special Forces Group (SFG) No. 26 CENEPA in Quevedo (Ecuador), which was subjected to a humid chamber to induce the manifestation of the causal agent(s) of the disease. This study area was selected based on the prevalence of disease symptoms in bamboo, which provides a relevant context for pathogen induction for the exact location of the study area (Figure 1). The experimental research was conducted in the "La María" Campus laboratories at the State Technical University of Quevedo, Ecuador.




Figure 1: Location of plant material.

Induction in a wet chamber

Plant material with signs of necrosis was placed in 14" x 20" polypropylene covers to induce the manifestation of disease-causing agents in bamboo. A controlled relative humidity was maintained for 72 to 96 hours. The effectiveness of this method was evaluated by observing morphological changes in the bamboo tissue, which were documented photographically. This technique is common in plant pathology to promote the manifestation of pathogens¹⁴.

Preparation of culture medium

Two modified culture media were prepared to promote the selective growth of fungi and bacteria. The PDA medium was enriched with Streptomycin (1 mg L^{-1}) to inhibit unwanted bacterial growth, while nutrient agar was used for bacterial cultivation. Both media were diluted in distilled water and autoclaved at 121°C for 25 minutes. These media were selected based on previous literature indicating their effectiveness in isolating pathogens from bamboo¹⁹.

Sowing in Petri plates

Aseptic samples of the induced plant material were seeded in Petri plates containing the prepared culture media²⁰. Pieces of bamboo culm were cut with a length and width of 2.5 cm and a thickness of 0.5 to 1 mm and incubated at 28 ± 2 °C for 8 days. Incubation was performed in a calibrated incubator to ensure consistency of growth conditions.

Sowing in carrot sandwiches

The carrot sandwich technique was used as a complementary method for pathogen induction. Carrots were washed, peeled, and disinfected with 80% ethanol before exposure to UV light for 15 minutes. The culm pieces were placed between two slices of carrot and secured with paper tape. This method allows a detailed observation of the pathogen-host interaction.

Isolation of microbial strains

Colonies developed in the Petri dishes were isolated using a sterile seed loop. They were transferred to new Petri containing the modified culture media [PDA (Streptomycin, 1mg L^{-1}) and nutrient agar (1mg L^{-1})] to obtain pure cultures²¹. This step was critical for the subsequent identification of the pathogens through biochemical and molecular tests

RESULTS

Description of the symptoms of the disease

The disease was observed to affect shoots in the early stages of development, with symptoms of internal rot evident in roots and young tissues at growth sections. This condition seriously compromises the production of new shoots, resulting in diminished natural regeneration, stunted growth and, ultimately, plant death. Affected shoots present dark brown to black exudates in the apical area, and the leaves are dark brown with black spots (Figure 2).




Figure 2: Comparison of healthy shoots and those affected by the disease. A) Healthy shoots from *D. asper*; B) *D. asper* shoots with exudations in the apical area.

The enzymatic action of *E. sinocalami* and *Arthrinium spp.* It resulted in spraying the internal part of the shoots, stopping the disease when reaching the culm area, which presents resistance due to its high cellulose content (Figure 3).




Figure 3: Evidence of enzymatic action on shoots and culm. The enzymatic action of *E. sinocalami* and *Arthrinium spp.* A) Internal part of the shoot pulverized by bacterial action; B) Attack by *E. sinocalami* and *Arthrinium spp.* in the culm inter-node.

Disease severity

The severity of the pathogen attack was reflected in a meager natural regeneration rate. The phenological evaluation showed that 97% of the shoots were affected and showed signs of spraying by phytopathogens (Table 1).

ITEM	Cane average	DCH	Live shoots (%)	Dead shoots (%)
Scale 1	0,2	8,80	3%	97%
Scale 2	10,0	12,50		
Scale 3	45,0	10,68		
Scale 4	3,0	14,16		
Scale 5	6,0	17,46		

Note: The phenological scale was arbitrarily based on vigor, diameter and hue.

Table 1. Phenological scale and percentage of affected shoots

Morphological recognition of microorganisms associated with bacterial wilt in *D. asper*

A strain of filamentous fungi and a bacterial strain of the genus *Erwinia*, closely related to bacterial wilt, were isolated and identified. Bacterial strains grew optimally on modified nutrient agar at room temperature, while fungal colonies preferred a PDA culture at 28°C. Identification was based on Gram staining and trypan blue pigmentation techniques, using microscopy with close-ups of 10, 40 and 100x and taxonomic keys to determine the genus of the strains.

Arthrinium species can produce varied fruiting bodies depending on growing conditions. It was observed that the aerial hyphae took on a white tone in the incubator and brown at room temperature. The identification of *Arthrinium* was confirmed by the observation of conidiogenous structures and spores with a germinal cleft (Figure 4).




Figure 4: Morphology of *Arthrinium spp.* Observed in culture. A) Macroscopic view of *Arthrinium spp.*; B) Spores with germinal cleft.

DISCUSSION

This study provides a detailed insight into the complex interaction between pathogenic microorganisms and *D. asper*, revealing the synergy between *E. sinocalami* and *Arthrinium spp.* in the induction of bacterial wilt and culm degradation. The complex interplay between microorganisms and bamboo, particularly in the context of disease management and control strategies, is a critical area of focus. Recent studies have shed light on various aspects of this interaction, offering insights that could be instrumental in developing more effective disease management strategies for *D. asper*. Darma et al.²² explored the potential of *Bacillus subtilis* BMB26, a bacterium isolated from bamboo powder, as a biocontrol agent against phytopathogenic fungi. This finding is particularly relevant to our study as it suggests the possibility of utilizing bamboo-associated microorganisms for biologically controlling pathogens in *D. asper*. The use of such biocontrol agents could provide a sustainable alternative to chemical control methods, aligning with the need for eco-friendly disease management practices in bamboo ecosystems²². The impact of forest management practices on the soil microbial community was highlighted by Yang et al.²³. Their study indicated that converting secondary broadleaf forests to Moso bamboo plantations alters the soil bacterial community structure. This alteration could have significant implications for disease dynamics in bamboo plantations, underscoring the importance of considering forest management practices in disease management strategies.

Zhang et al.²⁴ found that chicken farming under Moso bamboo forests significantly increased bacterial and fungal diversity. This increase in microbial diversity could influence the prevalence and severity of diseases such as bacterial wilt in bamboo, suggesting that agricultural practices in bamboo forests must be carefully managed to maintain ecological balance and prevent disease outbreaks²⁴.

Fuke et al.²⁵ emphasized rhizospheric microbial communities' role in bamboo health. These communities play a significant role in nutrient immobilization and phytoremediation, which could enhance the overall health and disease resistance of *Dendrocalamus asper*. Understanding and harnessing these microbial interactions could improve bamboo resilience against pathogens. The study by Shen et al.²⁶ on endophytic fungi isolated from Moso bamboo seeds revealed their broad-spectrum antimicrobial activity. These fungi could be a valuable source of bioactive compounds for developing plant defense activators, offering a novel approach to disease management in bamboo.

The interaction of fungi of the genus *Arthrinium* and bacteria *Erwinia spp.* with bamboo is complex and multifaceted. These microorganisms can exist in symbiosis with bamboo or become disease-causing pathogens²⁷. In particular, *E. sinocalami* and *Arthrinium spp.* have been identified as causing bacterial wilt and rot. of

shoots and culms, respectively²⁸. This study reveals a coordinated attack of these pathogens on bamboo shoots, suggesting more complex disease dynamics than previously recognized. The persistence of bacterial wilt in bamboo, associated with *E. sinocalami*, has been documented for over two decades²⁹. The limitation of this bacteria to attack only young shoots has significant implications for the regeneration and propagation of bamboo, as observed in the bamboo patches evaluated, where most shoots were affected or destroyed by pathogens. Identifying species of the genus *Arthrinium* is challenging due to the similarity of their conidia between different species and the morphological variability depending on the incubation environment^{30,31}. In this study, identification was achieved by observing conidiogenous structures and spores with a germinal cleft, which aligns with the existing literature's descriptions. Furthermore, it is essential to consider the role of natural products in crop protection, as discussed in the work of Dayan et al.³², which highlights the need to discover and develop new pesticides based on natural products to replace compounds lost due to stricter regulations. This could be relevant for the development of biological control strategies against pathogens in bamboo. The diversity of microorganisms in fermented foods and beverages globally, as reviewed in the study by Tamang et al.³³, also provides a context for understanding the complexity of microbial communities and their potential in biological control applications. Finally, the review by Bhardwaj et al.³⁴ on xylanases highlights the importance of these enzymes in the cleavage of complex polysaccharides and their application in various industrial and biotechnological sectors. Since *Arthrinium spp.* and *Erwinia spp.* are involved in the revision of plant tissues, understanding the enzymes involved could offer ways to mitigate their pathogenic impact. These findings and comparisons with existing literature underscore the need for a deeper understanding of interactions between pathogens and host plants. They could inform the development of integrated disease management strategies for the conservation and sustainable use of bamboo ecosystems.

CONCLUSIONS

This study provides a detailed insight into the complex interaction between pathogenic microorganisms and *D. asper*, revealing the synergy between *E. sinocalami* and *Arthrinium spp.* in the induction of bacterial wilt and culm degradation in bamboo. The findings demonstrate that *E. sinocalami* targets young shoots, while *Arthrinium spp.* is associated with the degradation of culm structure, suggesting a coordinated attack mechanism that could have significant implications on the health and viability of bamboo plantations. Identifying *Arthrinium spp.* was particularly challenging due to the morphological similarity of its conidia to other fungal species. However, accurate identification was achieved through detailed analysis of hyphae, conidiogenous structures and spores, highlighting the importance of meticulous diagnostic techniques in plant pathology. The results of this study emphasize the need to develop integrated disease management strategies that consider the interaction between different pathogens. Furthermore, it is crucial to continue research to understand better the epidemiology of bacterial wilt and pathogen-host interactions in bamboo to develop effective and sustainable control measures that ensure the protection of this important plant species in ecosystems: forestry and agricultural production. Collaboration between plant pathologists, microbiologists and agronomists will be essential to address the challenges presented by bacterial wilt in *D. asper* and other bamboo genera, ensuring the sustainability of these valuable resources worldwide. Finally, these studies collectively highlight the need for a holistic approach to disease management in bamboo ecosystems. By considering both beneficial and pathogenic microorganisms and the impact of environmental and management practices, we can develop more effective and sustainable strategies for the conservation and health of *D. asper*.

Author Contributions: Conceptualization, MV, JM; methodology, CB, MV, JM and MC; validation, JM, CB and RH; formal analysis JM, MV and CB; investigation, MV; CB and JM; writing-original draft

preparation, MV, JM, MC and RH; writing-review and edition, JM, MC, RH and MV; visualization, JM, MC, RH and MV; project administration, CB, MV and JM; funding acquisition, MV, CB and JM All authors have read and agreed to the published version of the manuscript.

Acknowledgments: The authors are grateful for the support of the Universidad Técnica Estatal de Quevedo (UTEQ).

Conflicts of Interest: There is no conflict of interest.

REFERENCES

1. Yuan, Z.-S. *et al.* Structural variability and differentiation of niches in the rhizosphere and endosphere bacterial microbiome of moso bamboo (*Phyllostachys edulis*). *Sci. Rep.* **11**, 1574 (2021).
2. Tu, Z., Chen, L., Yu, X. & Zheng, Y. Rhizosphere soil enzymatic and microbial activities in bamboo forests in southeastern China. *Soil Sci. plant Nutr.* **60**, 134–144 (2014).
3. Caycedo Lozano, L., Ramírez, L. C. C. & Suárez, D. M. T. Las bacterias, su nutrición y crecimiento: una mirada desde la química. *Nova* **19**, 49–94 (2021).
4. Cano, M. A. Interacción de microorganismos benéficos en plantas: Micorrizas, *Trichoderma* spp. y *Pseudomonas* spp. Una revisión. *Rev. UDCA Actual. Divulg. Científica* **14**, 15–31 (2011).
5. Morakotkarn, D., Kawasaki, H. & Seki, T. Molecular diversity of bamboo-associated fungi isolated from Japan. *FEMS Microbiol. Lett.* **266**, 10–19 (2007).
6. Tian, X.-K. *et al.* Native bamboo invasions into subtropical forests alter microbial communities in litter and soil. *Forests* **11**, 314 (2020).
7. Zhang, H. *et al.* Microbial taxa and functional genes shift in degraded soil with bacterial wilt. *Sci. Rep.* **7**, 39911 (2017).
8. Li, C., Zong, Z., Qie, H., Fang, Y. & Liu, Q. CiteSpace and Bibliometric Analysis of Published Research on Forest Ecosystem Services for the Period 2018–2022. *Land* **12**, 845 (2023).
9. Fujiwara, K., Aoyama, C., Takano, M. & Shinohara, M. Suppression of *Ralstonia solanacearum* bacterial wilt disease by an organic hydroponic system. *J. Gen. plant Pathol.* **78**, 217–220 (2012).
10. Li, S. *et al.* Soil acidification aggravates the occurrence of bacterial wilt in South China. *Front. Microbiol.* **8**, 703 (2017).
11. Santiago, T. R., Lopes, C. A., Caetano-Anollés, G. & Mizubuti, E. S. G. Phylotype and sequevar variability of *Ralstonia solanacearum* in Brazil, an ancient centre of diversity of the pathogen. *Plant Pathol.* **66**, 383–392 (2017).
12. Brooks, D. S., Gonzalez, C. F., Appel, D. N. & Filer, T. H. Evaluation of endophytic bacteria as potential biological-control agents for Oak Wilt. *Biol. Control* **4**, 373–381 (1994).
13. Kim, N. *et al.* Induction of resistance against pine wilt disease caused by *Bursaphelenchus xylophilus* using selected pine endophytic bacteria. *Plant Pathol.* **68**, 434–444 (2019).
14. Li, S.-J. *et al.* Purification of protein AP-Toxin from *Arthrinium phaeospermum* causing blight in *Bambusa pervariabilis* × *Dendrocalamopsis grandis* and its metabolic effects on four bamboo varieties. *Phytopathology* **103**, 135–145 (2013).
15. Dohet, L., Gregoire, J.-C., Berasategui, A., Kaltenpoth, M. & Biedermann, P. H. W. Bacterial and fungal symbionts of parasitic *Dendroctonus* bark beetles. *FEMS Microbiol. Ecol.* **92**, fiw129 (2016).
16. Lamichhane, J. R. & Venturi, V. Synergisms between microbial pathogens in plant disease complexes:

- a growing trend. *Front. Plant Sci.* **6**, 385 (2015).
17. van Dijk, L. J. A., Ehrlén, J. & Tack, A. J. M. The timing and asymmetry of plant–pathogen–insect interactions. *Proc. R. Soc. B* **287**, 20201303 (2020).
 18. Graziosi, I. & Rieske, L. K. A plant pathogen causes extensive mortality in an invasive insect herbivore. *Agric. For. Entomol.* **17**, 366–374 (2015).
 19. Mishra, R. K., Pandey, B. K., Muthukumar, M., Pathak, N. & Zeeshan, M. Detection of Fusarium wilt pathogens of *Psidium guajava* L. in soil using culture independent PCR (ciPCR). *Saudi J. Biol. Sci.* **20**, 51–56 (2013).
 20. Cheng, S.-F. *et al.* Identification of differentially expressed genes induced by bamboo mosaic virus infection in *Nicotiana benthamiana* by cDNA-amplified fragment length polymorphism. *BMC Plant Biol.* **10**, 1–12 (2010).
 21. Wang, H. *et al.* Dominance of zoonotic genotype D of *Enterocytozoon bieneusi* in bamboo rats (*Rhizomys sinensis*). *Infect. Genet. Evol.* **73**, 113–118 (2019).
 22. Darma, R. *et al.* A strong antifungal-producing bacteria from bamboo powder for biocontrol of *Sclerotium rolfsii* in melon (*Cucumis melo* var. *amanta*). *J. Plant Pathol. Microbiol.* **7**, 334 (2016).
 23. Yang, C., Zhong, Z., Zhang, X., Bian, F. & Du, X. Responses of soil organic carbon sequestration potential and bacterial community structure in moso bamboo plantations to different management strategies in subtropical China. *Forests* **9**, 657 (2018).
 24. Zhang, X. *et al.* Leaf-associated shifts in bacterial and fungal communities in response to chicken rearing under moso bamboo forests in subtropical China. *Forests* **10**, 216 (2019).
 25. Fuke, P., Kumar, M., Sawarkar, A. D., Pandey, A. & Singh, L. Role of microbial diversity to influence the growth and environmental remediation capacity of bamboo: a review. *Ind. Crops Prod.* **167**, 113567 (2021).
 26. Shen, X.-Y. *et al.* Diversity and antimicrobial activity of culturable endophytic fungi isolated from moso bamboo seeds. *PLoS One* **9**, e95838 (2014).
 27. Hyde, K. D., Zhou, D., McKenzie, E. H. C., Ho, W. H. & Dalisay, T. Vertical distribution of saprobic fungi on bamboo culms. *Fungal Divers.* (2002).
 28. Wang, S. *et al.* Electron transport chains in organohalide-respiring bacteria and bioremediation implications. *Biotechnol. Adv.* **36**, 1194–1206 (2018).
 29. Fonseca, W. & Rojas, M. Acumulación y predicción de biomasa y carbono en plantaciones de bambú en Costa Rica. *Ambient. y Desarro.* **20**, 85–98 (2016).
 30. Crous, P. W. & Groenewald, J. Z. A phylogenetic re-evaluation of *Arthrinium*. *IMA Fungus* **4** (1): 133–154. at (2013).
 31. Dai, D. Q. *et al.* Bambusicolous fungi. *Fungal Divers.* **82**, 1–105 (2017).
 32. Dayan, F. E., Cantrell, C. L. & Duke, S. O. Natural products in crop protection. *Bioorg. Med. Chem.* **17**, 4022–4034 (2009).
 33. Tamang, J. P., Watanabe, K. & Holzapfel, W. H. Diversity of microorganisms in global fermented foods and beverages. *Front. Microbiol.* **7**, 377 (2016).
 34. Bhardwaj, N., Kumar, B. & Verma, P. A detailed overview of xylanases: an emerging biomolecule for current and future prospective. *Bioresour. Bioprocess.* **6**, 1–36 (2019).

Received: October 9th 2023/ **Accepted:** January 15th 2024 / **Published:** 15 February 2024

Citation: Morante-Carriel J., Vera-Carranza M., Belezaca-Pinargote C., Carranza-Patiño M., Herrera-Feijoo R.J.. Microorganisms associated with bacterial wilt disease in *Dendrocalamus asper* (Giant Bamboo) from Ecuador. Revis Bionatura 2024; 9 (1) **32**. <http://dx.doi.org/10.21931/RB/2024.09.01.32>

Additional information Correspondence should be addressed to jmorante@uteq.edu.ec

Peer review information. Bionatura thanks anonymous reviewer(s) for their contribution to the peer review of this work using <https://reviewerlocator.webofscience.com/>

All articles published by Bionatura Journal are made freely and permanently accessible online immediately upon publication, without subscription charges or registration barriers.

Bionatura ISSN. First 13909355 Ecuador. **Scopus coverage years:** from 2016 to the present

Publisher's Note: Bionatura stays neutral concerning jurisdictional claims in published maps and institutional affiliations.

Copyright: © 2023 by the authors. They were submitted for possible open-access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).