

Article

Pathology and immunological changes of Typhoid fever

Rima N. Hasan¹ and Ali Anok Njum^{2,*}

¹ Department of Community health, Techniques, Al-Furat Al-Awsat Technical University , Technical Institute/ Samawa;

² Department of Community health, Techniques, Al-Furat Al-Awsat Technical University , Technical Institute/ Samawa;

* Correspondence: Aliscience16@yahoo.com

Abstract: A total of 150 blood and stool samples maturing between the ages of 10 and 60 were collected and cultured on BHI medium, MacConkey, and XLD. agars were all found to contain *Salmonella typhi* after biochemical testing for the bacteria. Disconnects of *Salmonella typhi* and 12 segregates of the pathogen were detected in the stool and blood cultures, respectively. Similarly, seropositive serum samples from patients infected with salmonella tested by ELISA assay to evaluate the concentration of cytokines and immune marks (IL-18, TNF-B, CD8 and CD4). Results showed that acute cases of disease express high level of cytokine and immune marks as compared with chronic and asymptomatic infection

Keyword: Immunity; CD Markers; cytokines; salmonella.

1. Introduction

Salmonella is the most frequent genus in the Enterobacteriaceae family. Bacteria that are Gram-ve facultative anaerobes may thrive in a wide range of habitats and hosts, including humans, animals, as well as food. For example, *Salmonella* variety Humans and other domesticated or wild species may be infected with *Salmonella* bacterium¹. It's a common theory that this food-borne illness-causing bacteria is to blame². These microscopic organisms are to blame for disorders like enteritis and enteric fever^{3,4}. Anywhere from three days to nearly a month ago, the illness first surfaced, and the death rate might be anywhere from (12 percent and 30 percent). While typhoid fever's total impact is reduced, there has been a noticeable increase in the number of cases. Cytokines grind as the particles of sentinel retort that convey about several functional occupations and variation the defending, Lymphocyte cytokines are primarily involved in the host response^{5,6}. Invigorated lymphocytes are divided into two classes that release cytokines. T helper 1 cytokines, such as IL-12 and IFN-, stimulate the (CMI) response, whereas T helper 2 cytokines, such as IL-8 and IL-18, are concerned about AMI. The two reactions have been discovered to be linked in a viral infection^{7,8} and the differences between them tilt toward HIR and discouraged alter CMI, which is essential for infection resistance^{9,10}

2. Materials and methods

At the emergency clinic, a doctor recorded clinical indications of typhoid tolerance and immediately determined to conduct a Widal test on all 150 patients. Stool samples were cultured directly on macConky agar ,while blood cultured on blood culture medium then isolated on macConky and XLD agar, then identified by biochemical tests and vitek system.

2.1. Serological test

patient blood was drained for each of patients. The serum from the blood tests was centrifuged, and the Widal test and IgG and IgM strips were used to evaluate the typhoid patients' seropositive. levels of cytokines and CD8 and CD4 immunomarkers using the ELISA methods

2.2. Statistical analysis

It was exhibited by expending Chi-square (χ^2) examination to control the numerical variations between various collections by consuming an application statistical stand for social science (SPSS 19). The opportunity of ($P \leq 0.05$) was restrained to be statistically important.

3. Results

3.1. Clinical sings

In a sample of 150 "blood and stool samples" from persons with suspected typhoid fever, six disengages 12 isolates of *S. Typhi* were found. Death and desolation are common consequences of typhoid disease. In Iraq, screenings for typhoid fever were made mandatory. Typhoid fever

outbreaks have been researched extensively in these investigations; they discovered that cases spread across a large area inside the league, increasing the danger of water-borne illness¹¹ according serological test showed only 50 patients (33.33%) were seropositive for salmonella and 30 patient showed chronic infection as well as 20 patient showed acute disease this results as showed by(12)Serum of all patients with typhoid and those with intense or persistent illness activity contain more elevated level of IL-18, TNF-B, CD2 and CD4 than chronic disease as compared with control groups .as tables(1,2,3 and 4)

Table 1. The Concentration of IL-18.

Group	NO.	Serum level of IL-18		
		Mean	low	high
Acute	20	1000	500	1100
chronic	30	700	400	900
Control	20	10	5	11

Table 2. The Concentration of TNF-B.

Group	NO.	Serum level of TNF-B		
		Mean	low	high
Acute	20	600	500	800
chronic	30	150	50	200
Control	20	12	5	20

Table 3. The appearance of CD8

Group	NO.	Serum level of CD8		
		Mean	low	high
chronic	30	5	2	9
Acute	20	15	8	19
Control	20	3	1	4

Table 4. The Concentration of CD4

Group	NO.	Serum level of CD4		
		Mean	low	high
Acute	20	30	19	38
chronic	30	15	8	17
Control	20	2	1	4



Figure 1. show Salmonella typhi culture

4. Discussion

Chemokines work by connecting with certain cell surface receptors on the body. Inconsistencies within the structure of chemokine receptors account for their extraordinary attraction to a wide variety of ligands^{13,14}. Neutrophils and monocytes are reactivated by IL-8 through inspiration as well as induction^{1,16}. Neutrophils serve as the initial line of defense rather than fighting a wide variety of germs as they would in an infection condition. Provocative cytokines, including IL-8, IL-10, and IL-12, are released by these cells, which also create irritating oxygen. The release of IL-8 into the lungs is associated with an increase in neutrophil activity^{11,7}. The beginning of IL-8 can be expanded. Aside from this, the release of responsive O₂-species from granulated cells was found to disrupt the appearance of "IL-18." In patients with severe S.typhi pollution, lung IL-18 is seen at a lower level of protection, but blood serum and liver examination can be used to identify individuals with moderate disease. CD4⁺ and CD8⁺ White blood cell reactions associated with the removal of S.typhi are critical for those who have a mild infection to have delicate, slimly focused reactions^{20,19}. CD8⁺ effector cells in the lung began to lose their ability to operate, as seen by a decrease in IFN- γ production. The protection of lung microorganisms is as frequently as feasible provided by fragile antigens resulting from the "CD8⁺ White blood cell reaction." Due to the difficulty in determining whether CD8 was pathogenic or not, our results show that both CD4 and CD8 are up-regulated in the peripheral blood of Coronavirus patients in a state of resistant dysregulation.^{22,21}

5. Conclusion

Salmonella is the most frequent genus in the Enterobacteriaceae family. Bacteria that are Gram-negative facultative anaerobes may thrive in a wide range of habitats and hosts, including humans, animals, as well as food. The sample of 150 "blood and stool samples" from persons with suspected typhoid fever, showed that acute cases of disease express high level of cytokine and immune marks as compared with chronic and asymptomatic infection. The protection of lung microorganisms is as frequently as feasible provided by fragile antigens resulting from the "CD8⁺ White blood cell reaction." Due to the difficulty in determining whether CD8 was pathogenic or not, our results show that both CD4 and CD8 are up-regulated in the peripheral blood of Coronavirus patients in a state of resistant dysregulation.

References

- Kariuki, S.; Revathi, G.; Kariuki, N.; Kiiru, J.; Hart, C.A. Characterisation of community acquired non-typhoidal Salmonella from bacteraemia and diarrhoeal infections in children admitted to hospital in Nairobi, Kenya. *BMC microbiology*, 2006, 6.1: 1-10.
- Kaseb, A. O.; Chinnakannu, K.; Chen, D.; Menon, M.; Dou, Q. P. Androgen receptor- and E2F-1-targeted thymoquinone therapy for hormone-refractory prostate cancer. *Cancer research*, 2007, 67.16: 7782-7788.
- Kayashima, S.; Tsuru, S.; Shinomiya, N.; Katsura, Y.; Motoyoshi, K.; Rokutanda, M.; Nagata, N. Effects of macrophage colony-stimulating factor on reduction of viable bacteria and survival of mice during *Listeria monocytogenes* infection: characteristics of monocyte subpopulations. *Infection and immunity*, 1991, 59.12: 4677-4680.
- Khan, M. I.; Fadl, A. A.; Venkitanarayanan, K. S. Reducing colonization of *Salmonella* Enteritidis in chicken by targeting outer membrane proteins. *Journal of Applied Microbiology*, 2003, 95.1: 142-145.
- Khather, Z. K.; Shareef, A. M.. Contamination of the parentstock feed with salmonella in Ninawa Government. *Iraqi Journal of Veterinary Sciences*, 19(2), 85-95.
- Kilic, A.; Bay Sallar, M.; Salih, B.; Sorkun, K.; Tanyuksel, M. . In vitro antimicrobial activity of propolis against methicillin-resistant *Staphylococcus aureus* and vancomycin-resistant *Enterococcus faecium*. *Annals of Microbiology*, 2005, 55.2.
- Tizard, I.R. An introduction to veterinary immunology. WB Saunders., 1977.
- Torpdahl, M.; Sorensen, G.; Lindstedt, B.A. Tandem repeat analysis for surveillance of human *Salmonella* Typhimurium infections. *Emerging infectious diseases*, 2007, 13.3: 388.
- Trent, M.S.; Stead, C.M.; Tran A.X.; Hankins, J.V. Diversity of endotoxin and its impact on pathogenesis. *J. Endotoxin. Res.*2006. 12: 205-23.
- Uchiya, K.; Barbieri, M. A.; Funato, K. A. A *Salmonella* virulence protein that inhibits cellular trafficking. *The EMBO journal*, 1999, 18.14: 3924-3933.
- Van Duijkereen, E. W. J. B.; Wannet, W. J. B.; Houwers, D. J.; Van Pelt, W. Serotype and phage type distribution of *Salmonella* strains isolated from humans, cattle, pigs, and chickens in the Netherlands from 1984 to 2001. *Journal of Clinical Microbiology*, 2002, 40.11: 3980-3985.
- Vandepitte, L.; Engbac, K.; Pito, P.; Hench, C. C. Basic laboratory procedures in clinical bacteriology. World Health Organization, 2003.
- Suwaid, A. H. ; Rashid, M. A. ; Taha, M. M. . Genetic Analysis For Combining Ability And Estimation Of Some Genetic Parameters Of Yield And Its Components In Maize Using Half Diallel Cross.). *Journal of Life Science and Applied Research*. 2020, 1, 60-64.
- Vazquez-Torres, A.; Fang, F. C. Cellular routes of invasion by enteropathogens. *Current opinion in microbiology*, 2000, 3.1: 54-59.
- Vetvicka, V.; Vetvickova, J. An evaluation of the immunological activities of commercially available β 1, 3-glucans. *JANA*, 2007, 10.1: 9-15.
- Noaman A I, Khalaf R M, Emad GH, Al-Abbasy, Mohammed Th. T. Effect of flaxseed oil dosing on fertility, growth characteristics and some physical, biochemical, and hormonal blood parameters during the early pregnancy of Awassi ewes. *Revis Bionatura*. 2022;7(4) 5. <http://dx.doi.org/10.21931/RB/2022.07.04.5>.
- Vogal, F.R.; Powll, M.F.; Michael, F. A compendium of vaccine adjuvants and excipients. *Vaccine Design*, 1995, 141-228..
- Wain, J.; House, D.; Baker, S.; Kidgell, C.; Rumina, H. Vi antigen expression in *Salmonella enterica* serovar Typhi clinical isolates from Pakistan. *Journal of clinical microbiology*, 2005, 43.3: 1158-1165.

19. Waterman, S. R.; Holden, D.W. Identification of salmonella pathogenicity island-2 type III secretion system effectors involved in intra-macrophage replication of *S. enterica* serovar typhimurium: implications for rational vaccine design. *MBio*, 2013, 4.2: e00065-13.
20. Weill, F.X.; Timinouni, M.; Demartin, M.; Polomack, L.; Grimont, P. A. Multi-drug resistance to *Salmonella typhimurium* from human influenza. *J.Clin.Microbio.* 2006, 44 (3): 701-708.
21. Control and Prevention diseases centre (CDC). Outbreak of *Salmonella* serotype Saintpaul infections associated with multiple raw produce items--United States, 2008. *MMWR. Morbidity and mortality weekly report*, 2008, 57.34: 929-934.
22. Control and Prevention diseases centre (CDC). Preliminary FoodNet data on the incidence of infection with pathogens transmitted commonly through food--selected sites, United States, 2003. *MMWR. Morbidity and mortality weekly report*, 2004, 53.16: 338-343